

From: BuyandBill.com <info@buyandbill.com>
Sent: XXXXX
To: XXXXXX
Subject: Announcing a price change for Paragard®



CooperSurgical®

1-877-PARAGARD

Paragard Customer,

Effective September 30, 2019, the pricing of Paragard will change as outlined below.

Please take a moment to update any of your reimbursement, billing, and ordering systems to reflect this change.

PARAGARD NDC CODES:

HCPCS Code	Labeler	11 Digit NDC	NDC Effective Date	Product Description	WAC	Estimated AWP
J7300	COOPERSURGICAL	59365-5128-01	9/30/19	Carton of 1 (one) sterile Unit	\$884.50	\$1061.40
J7300	TEVA WOMENS HEALTH	51285-0204-01	9/30/19	Carton of 1 (one) sterile Unit	\$884.50	\$1061.40

CooperSurgical is notifying all customers and business associates about this change and we would appreciate your partnership in helping us ensure that all systems and locations involved with Paragard are updated with the new Paragard pricing.

For any questions, contact CooperSurgical at 1.877.PARAGARD.

Paragard is a copper-releasing intrauterine device (IUD) that is used to prevent pregnancy.

- 100% hormone free and over 99% effective
- Lasts for up to 10 years, but can be removed at any time
- Immediately reversible; women can start trying to get pregnant the same day it's removed

Indication

Paragard is indicated for intrauterine contraception for up to 10 years.

Important Safety Information

- Paragard must not be used by women who have acute pelvic inflammatory disease (PID); have had a postpregnancy or postabortion uterine infection in the past 3 months; have cancer of the uterus or cervix; have an infection of the cervix; have an allergy to any component; or have Wilson's disease.
- If a woman misses her period, she must be promptly evaluated for pregnancy.
- Possible serious complications that have been associated with intrauterine contraceptives are PID, embedment, perforation of the uterus, and expulsion.
- Paragard must not be used by women who are pregnant as this can be life threatening and may result in loss of pregnancy or infertility.
- The most common side effects of Paragard are bleeding and spotting; for most women, these typically subside after 2 to 3 months.
- **Paragard does not protect against HIV or other sexually transmitted infections (STI).**

Click here for the full [Prescribing Information](#) for Paragard.

US-PAR-1900140 **September 2019**

buyandbill.com, 30 Fawn Run, Glastonbury, CT 06033

[SafeUnsubscribe™ {recipient's email}](#)

[Forward this email](#) | [Update Profile](#) | [About our service provider](#)

Sent by info@buyandbill.com in collaboration with

Constant Contact 

[Try email marketing for free today!](#)